

For patients 12 and older, after 2 chronic GVHD systemic treatments haven't worked...

Open your world to possibilities with REZUROCK

See how REZUROCK may help
improve your chronic GVHD.

REZUROCK[®]
(belumosudil) tablets



Patient portrayal.
All individuals shown are actors.

GVHD, graft-versus-host disease.

USE

REZUROCK[®] (belumosudil) is a prescription medicine used to treat adults and children 12 years of age and older with chronic graft-versus-host disease (chronic GVHD) after you have received at least 2 prior treatments (systemic therapy) and they did not work. It is not known if REZUROCK is safe and effective in children less than 12 years old.

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney or liver problems.
- are pregnant or plan to become pregnant. REZUROCK can harm your unborn baby. If you are able to become pregnant, your healthcare provider will do a pregnancy test before starting treatment with REZUROCK. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with REZUROCK.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

Taking control of your chronic GVHD journey

When chronic GVHD develops, it causes your immune system to become unbalanced, creating too many of some cells and molecules, and not enough of others, which can harm different parts of your body.

The imbalance in the immune system caused by chronic GVHD can lead to inflammation (swelling) and fibrosis (hardening and scarring), which can damage your organs.

Inflammation and fibrosis: the two challenges of chronic GVHD



Inflammation is swelling that occurs when your body is trying to protect itself.



Fibrosis is the growth of scar tissue that hardens and keeps your organs from working properly. In some people, fibrosis happens when inflammation continues for a long time. It is also possible for scar tissue to form without any inflammation.

To match you with a treatment for chronic GVHD, your health care team may consider several important factors, including:



The overall severity of your chronic GVHD



The effect chronic GVHD has on your daily life and activities



Which organs are involved

In some cases, your doctor may also recommend local treatments to help manage specific issues, for example, eye drops for eye dryness or creams for skin rashes. These may be used as part of your overall treatment plan.

Taking control of your chronic GVHD journey (cont)



Medicines that treat chronic GVHD

When it comes to treating chronic GVHD, there are different types of treatments available. Most patients start on an immunosuppressive therapy to help control their symptoms. Some people may need a different approach. Chronic GVHD is a long-term disease, and treatments may stop working over time.



Chronic GVHD medicines can help:

- Relieve symptoms
- Address the multiple impacts chronic GVHD can have on the body



Remember, you have the power to advocate for your treatment plan

If you are still noticing symptoms or feel like your medicine isn't working as well as it used to, or your condition isn't as well controlled as before, talk with your health care team.

With your health care team, you can explore if it may be time for other options that could be a better fit for you.

You know your body best.

If you notice any changes in how you're feeling, write them down and share with your health care team during your next appointment. This can help you understand what these symptoms might mean and whether your current treatment is still meeting your needs.



Click or Tap the QR code to download a tool that can help you keep track of your symptoms and aid conversations with your health care team.

An overactive ROCK2 pathway may play a role in chronic GVHD. How does this impact your body?




What is the ROCK2 pathway?

Many systems in the body work through what are called "pathways." These pathways are a series of actions that take place among cells and molecules in the body. One such pathway is called "ROCK2." In people with chronic GVHD, this ROCK2 pathway is too active.

An overactive ROCK2 pathway leads to:

- A decrease in cells and molecules that help keep the immune system in check
- An increase in cells and molecules that cause inflammation and fibrosis



The result of an overactive ROCK2 pathway is an immune system that is out of balance.

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you: (cont)

- **Females** who can become pregnant should use effective birth control during treatment with REZUROCK and for 1 week after the last dose.
- **Males** with female partners who can become pregnant should use effective birth control during treatment with REZUROCK and for 1 week after the last dose.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

Open your world to a different way of treating chronic GVHD




What is REZUROCK?

REZUROCK is a once-daily tablet for patients 12 years and older with chronic GVHD after 2 prior systemic treatments did not work.

How does REZUROCK work?

REZUROCK works differently by blocking the ROCK2 pathway. This may reduce the inflammation and fibrosis that can happen with chronic GVHD.



REZUROCK can help rebalance your immune system.



Open your world to how REZUROCK works.
Click or Tap the QR code to watch a video of REZUROCK in action.

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you: (cont)

- are breastfeeding or plan to breastfeed. It is not known if REZUROCK passes into breast milk. Do not breastfeed during treatment with REZUROCK and for 1 week after the last dose.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

Understanding the REZUROCK clinical studies

REZUROCK[®]
(belumosudil) tablets

REZUROCK is for patients 12 years and older with chronic GVHD after 2 prior systemic treatments did not work

REZUROCK was evaluated in:

2 clinical studies

Included a total of:

186 people with chronic GVHD

The ROCKstar study had:

132 people

The KD025-208 study had:

54 people

IMPORTANT SAFETY INFORMATION

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REZUROCK may affect the way other medicines work, and other medicines may affect the way REZUROCK works. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

Jen

Real REZUROCK
patient
compensated
by Sanofi.



Different groups of people took REZUROCK in the studies

REZUROCK[®]
(belumosudil) tablets

People who were evaluated in the clinical studies included:



People who tried immunosuppressants, such as steroids, and other medicines for chronic GVHD that failed to control their symptoms



People with mild, moderate and severe chronic GVHD



People with chronic GVHD in 1 or more organs



There's more to learn about the results from the **ROCKstar** study.

Click or Tap the QR code to visit [REZUROCK.com](https://www.rezurock.com).



Jen, real REZUROCK patient compensated by Sanofi. Other individual shown is an actor.

IMPORTANT SAFETY INFORMATION


How should I take REZUROCK?

- Take REZUROCK exactly as your healthcare provider tells you to take it.
- Do not change your dose or stop taking REZUROCK without first talking to your healthcare provider.
- Take REZUROCK 1 time a day with a meal.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

How REZUROCK may help improve your chronic GVHD

REZUROCK[®]
(belumosudil) tablets



3 out of 4 people

75% of people in the ROCKstar study saw improvement in their chronic GVHD

- More than half of people who saw improvement did so between **weeks 4 and 8**
- Almost everyone who saw improvement did so by **week 24**
- These results are based on clinical study data on file. Medicine(s) may affect everyone differently. Not everyone will respond and some people may take longer to respond, especially if they have severe disease caused by fibrosis. Individual results may vary.

IMPORTANT SAFETY INFORMATION

How should I take REZUROCK? (cont)

- Take REZUROCK at about the same time each day.
- Swallow REZUROCK tablets whole with a glass of water.
- Do not cut, crush, or chew REZUROCK tablets.
- Your healthcare provider will do blood tests to check your liver at least 1 time a month during treatment with REZUROCK.


Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).



Patient portrayal.

Responses were seen in various organs



Some people taking REZUROCK had:

- ✓ An observed **reduction in steroid use**^a
- ✓ **Quality of life improvements**, reported in an exploratory analysis^b

^aNo definitive conclusions about the effect of REZUROCK can be drawn from these data.

^bAn exploratory analysis provides a better understanding of a problem, but is not proof. It is difficult to measure quality of life because people are different and will have unique experiences with treatment.

No definitive conclusions about the effect of REZUROCK can be drawn from this data. Also, the ROCKstar study measured overall response across all organs, so improvement in one organ may not mean overall benefit.

IMPORTANT SAFETY INFORMATION

How should I take REZUROCK? (cont)

- If you miss a dose of REZUROCK, take it as soon as you remember on the same day. Take your next dose of REZUROCK at your regular time on the next day. Do not take extra doses of REZUROCK to make up for a missed dose.
- If you take too much REZUROCK, call your healthcare provider or go to the nearest hospital emergency room right away.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

What are the side effects of REZUROCK?

These were the most common side effects reported by people taking REZUROCK in the 2 clinical studies

- Infections
- Tiredness or weakness
- Nausea
- Diarrhea
- Shortness of breath
- Cough
- Swelling
- Bleeding
- Stomach (abdominal) pain
- Muscle or bone pain
- Headache
- High blood pressure

REZUROCK may affect fertility in males and females. Talk with your health care provider if this is a concern for you.

These are not all the possible side effects of REZUROCK.

Additional information from the clinical studies



In clinical studies, there were no new cases of cytomegalovirus (CMV) infection and only 1 case of recurring CMV infection. CMV is a common serious infection that can happen in people who have had a blood stem cell transplant.



About 4% of people, or 1 out of every 25, in clinical studies had serious cytopenia (SY-toh-PEE-nee-uh), a condition in which there is a lower-than-normal level of certain blood cells.

REZUROCK side effects (cont)



If you have chronic GVHD, talk with your health care provider about staying on REZUROCK as long as needed.

This is very important. Be sure to tell your health care provider about any side effects you may have. They may be able to help you find ways to manage them.

Patient portrayal.


IMPORTANT SAFETY INFORMATION

What are the possible side effects of REZUROCK?

The most common side effects of REZUROCK include:

- infections
- tiredness or weakness
- nausea
- diarrhea
- shortness of breath
- cough
- swelling
- bleeding
- stomach (abdominal) pain
- muscle or bone pain
- headache
- high blood pressure

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).



How to take REZUROCK



The REZUROCK 200-mg tablet is taken by mouth once a day with a meal



How and when do I take REZUROCK?

Take REZUROCK at approximately the same time each day with a meal. Remember, a meal should be similar in calories to what you would eat for lunch or dinner—it's more than just a snack.



Can I crush or chew the REZUROCK tablet?

The REZUROCK tablet should be swallowed whole with a glass of water. It is very important that you **do not cut, crush or chew the tablet**.



What happens if I miss a dose of REZUROCK?

If you miss a dose of REZUROCK, take it as soon as you remember on the same day. Take your next dose of REZUROCK at your regular time on the next day.

Do not take an extra dose of REZUROCK to make up for a missed dose.

IMPORTANT SAFETY INFORMATION

Your healthcare provider may change your dose of REZUROCK, temporarily stop, or permanently stop treatment with REZUROCK if you have certain side effects.

REZUROCK may affect fertility in males and females. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects of REZUROCK. Call your doctor for medical advice about side effects.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).



How do I store REZUROCK?

REZUROCK should be stored at room temperature, ranging from 68 °F–77 °F (20 °C–25 °C) in its original container to protect it from moisture. Make sure to close the cap tightly each time after taking your tablet.

Do not discard the drying packet inside the bottle.



How to take REZUROCK (cont)

Be sure to take once-daily REZUROCK as prescribed by your health care team

- Always talk with your health care team about any changes to your medications. Know the medicines you take. Keep a list of them to show your health care provider and pharmacist when you get a new medicine
- Do not start or stop taking any medicines without discussing it first with your health care team. If you experience a side effect, be sure to tell your health care team right away
- If you are taking REZUROCK, make sure to tell your health care team about all of the prescription and over-the-counter medicines (including proton pump inhibitors, or PPIs), vitamins and herbal products you are taking because they may affect how REZUROCK works. REZUROCK may affect the way other medicines work



Click or Tap the QR code

to hear from real patients and their experience with taking REZUROCK.

IMPORTANT SAFETY INFORMATION

Your healthcare provider may change your dose of REZUROCK, temporarily stop, or permanently stop treatment with REZUROCK if you have certain side effects. (cont)

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call **1-800-FDA-1088**. You may also contact Kadmon Pharmaceuticals, LLC, a Sanofi Company, at 1-800-633-1610 to report side effects.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

Jen

Real REZUROCK patient compensated by Sanofi. Other individuals shown are actors.



MyROCK ASSIST™ is here to help

REZUROCK®
(belumosudil) tablets

MyROCK
ASSIST™

**MyROCK ASSIST strives to help people with chronic GVHD
and their caregivers get what they need, every step of the way**

The MyROCK ASSIST team is available to help you
Monday through Friday, 8 AM-8 PM ET.

**MyROCK ASSIST is available to
help eligible people:**

- ✓ Determine insurance coverage and costs for REZUROCK
- ✓ Find a savings program they may qualify for
- ✓ Have medicine delivered directly by a specialty pharmacy

Sanofi does not provide medical advice, diagnosis or
treatment recommendations.

Patient portrayal.
All individuals shown are actors.



MyROCK ASSIST™ is here to help



Eligible commercially or privately insured people can receive REZUROCK for AS LOW AS \$0 PER MONTH through the MyROCK ASSIST Commercial Co-pay Savings Program.^a

Learn more about the MyROCK ASSIST support programs designed for people who need treatment for chronic GVHD at **REZUROCK.com**, or call **1-844-523-6661**.



Click or Tap the QR code to enroll today.



Patient portrayal. All individuals shown are actors.

^aPatient Terms and Conditions: The MyROCK ASSIST Commercial Co-pay Savings Program provides co-pay/coinsurance support for out-of-pocket costs on REZUROCK® (belumosudil) tablets prescriptions. A yearly maximum benefit applies. Limit one 30-day supply per 30 days. This program is not health insurance. This program is for commercially or privately insured patients only; uninsured or cash-paying patients are not eligible. Patients are not eligible if prescriptions are paid, in whole or in part, by any state- or federally funded programs, including, but not limited to, Medicare (including Part D, even in the coverage gap) or Medicaid, Medigap, VA, DOD, TriCare, private indemnity or HMO insurance plans that reimburse you for the entire cost of your prescription drugs, or where prohibited by law. The co-pay program may not be combined with any other rebate, coupon or offer. Sanofi reserves the right to rescind, revoke or amend this offer at any time without further notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. This program is intended to help patients afford REZUROCK. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Card is valid through December 31 of the year of activation. On January 1 of the following year, the card automatically resets and is subject to annual limits if the prescription benefit remains the same. A representative of Sanofi may contact the patient for follow up on any adverse event that may be reported. Upon registration, patients receive all program details.

Preparing for your conversation about REZUROCK



The time you have with your health care team is important. Preparing questions in advance helps ensure you make the most of your visit. You may use this guide to note questions about REZUROCK. Along with this guide, bring any other trackers or a journal you may have to help with this conversation.

During your visit:



Advocate for yourself:

If you have questions about your current medicine or treatment plan, now is a good time to ask your doctor.



Update your doctor on anything new:

Symptoms can affect everyone differently, so speaking from your own experience can help your doctor understand what you are facing.

To help prepare you for your next appointment, write your top 3 questions about REZUROCK:

1. _____
2. _____
3. _____

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney or liver problems.
- are pregnant or plan to become pregnant. REZUROCK can harm your unborn baby. If you are able to become pregnant, your healthcare provider will do a pregnancy test before starting treatment with REZUROCK. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with REZUROCK.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

Use and Important Safety Information



USE

REZUROCK® (belumosudil) is a prescription medicine used to treat adults and children 12 years of age and older with chronic graft-versus-host disease (chronic GVHD) after you have received at least 2 prior treatments (systemic therapy) and they did not work. It is not known if REZUROCK is safe and effective in children less than 12 years old.

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney or liver problems.
- are pregnant or plan to become pregnant. REZUROCK can harm your unborn baby. If you are able to become pregnant, your healthcare provider will do a pregnancy test before starting treatment with REZUROCK. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with REZUROCK.
 - **Females** who can become pregnant should use effective birth control during treatment with REZUROCK and for 1 week after the last dose.
 - **Males** with female partners who can become pregnant should use effective birth control during treatment with REZUROCK and for 1 week after the last dose.
- are breastfeeding or plan to breastfeed. It is not known if REZUROCK passes into breast milk. Do not breastfeed during treatment with REZUROCK and for 1 week after the last dose.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REZUROCK may affect the way other medicines work, and other medicines may affect the way REZUROCK works.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I take REZUROCK?

- Take REZUROCK exactly as your healthcare provider tells you to take it.
- Do not change your dose or stop taking REZUROCK without first talking to your healthcare provider.
- Take REZUROCK 1 time a day with a meal.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

Use and Important Safety Information (cont)



IMPORTANT SAFETY INFORMATION

How should I take REZUROCK? (cont)

- Take REZUROCK at about the same time each day.
- Swallow REZUROCK tablets whole with a glass of water.
- Do not cut, crush, or chew REZUROCK tablets.
- Your healthcare provider will do blood tests to check your liver at least 1 time a month during treatment with REZUROCK.
- If you miss a dose of REZUROCK, take it as soon as you remember on the same day. Take your next dose of REZUROCK at your regular time on the next day. Do not take extra doses of REZUROCK to make up for a missed dose.
- If you take too much REZUROCK, call your healthcare provider or go to the nearest hospital emergency room right away.

What are the possible side effects of REZUROCK?

The most common side effects of REZUROCK include:

- | | |
|-------------------------|----------------------------|
| • infections | • swelling |
| • tiredness or weakness | • bleeding |
| • nausea | • stomach (abdominal) pain |
| • diarrhea | • muscle or bone pain |
| • shortness of breath | • headache |
| • cough | • high blood pressure |

Your healthcare provider may change your dose of REZUROCK, temporarily stop, or permanently stop treatment with REZUROCK if you have certain side effects.

REZUROCK may affect fertility in males and females. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects of REZUROCK. Call your doctor for medical advice about side effects.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call **1-800-FDA-1088**. You may also contact Kadmon Pharmaceuticals, LLC, a Sanofi Company, at 1-800-633-1610 to report side effects.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

For patients 12 and older, after 2 chronic GVHD systemic treatments haven't worked...

Open your world to possibilities with REZUROCK

See how REZUROCK may help improve your chronic GVHD

- REZUROCK is the only approved medicine that blocks the ROCK2 pathway, which could help **rebalance** your immune system. It **may reduce inflammation and fibrosis** that can happen with chronic GVHD
- **75% of people** (that's 3 out of every 4 people) in the ROCKstar study saw **improvement in their chronic GVHD**
- More than half of people who saw improvement did so between **weeks 4 and 8**^a
- Some people had an **observed reduction in steroid use**^b
- **Quality of life improvements was seen by some people**, reported in an exploratory analysis^c
- REZUROCK is a once-daily oral **tablet**, taken with a meal at the **same time each day**
- **Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including** if you have kidney or liver problems, are pregnant or planning to become pregnant, or are breastfeeding or plan to breastfeed. Both females who can become pregnant and males with female partners who can become pregnant should use effective birth control during treatment and for one week after the last dose
- **The most common side effects** of REZUROCK include infections, tiredness or weakness, nausea, diarrhea, shortness of breath, cough, swelling, bleeding, stomach (abdominal) pain, muscle or bone pain, headache, and high blood pressure



Click or Tap the QR code or visit REZUROCK.com, where you'll find everything you need about chronic GVHD, REZUROCK and MyROCK ASSIST™.

^aThese results are based on clinical study data on file. Individual results may vary.

^bNo definitive conclusions about the effect of REZUROCK can be drawn from these data.

^cAn exploratory analysis provides a better understanding of a problem, but is not proof. It is difficult to measure quality of life because people are different and will have unique experiences with treatment.

USE

REZUROCK® (belumosudil) is a prescription medicine used to treat adults and children 12 years of age and older with chronic graft-versus-host disease (chronic GVHD) after you have received at least 2 prior treatments (systemic therapy) and they did not work. It is not known if REZUROCK is safe and effective in children less than 12 years old.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).

Sanofi does not provide medical advice, diagnosis or treatment. The health information contained herein is provided for general educational purposes only. Your health care professional is the best source of information regarding your health. Please consult your health care professional if you have any questions about your health or treatment.

sanofi

© 2026 Sanofi. All rights reserved.
All trademarks mentioned are the property of the Sanofi group.
MAT-US-2203537-v6.0-06/2026

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you: (cont)

- **Females** who can become pregnant should use effective birth control during treatment with REZUROCK and for 1 week after the last dose.
- **Males** with female partners who can become pregnant should use effective birth control during treatment with REZUROCK and for 1 week after the last dose.

REZUROCK®
(belumosudil) tablets